

Find the maximum element of array
A[0..n-1] by iteratively setting A[i] to
 $\max(A[2i], A[2i+1])$

T(n) = _____ T(_____) + _____

Solving Recurrence Equations: The Iteration Method


The **iteration method** is a two-step strategy for solving recurrence equations:

1. Iteratively construct a recursion tree by unwinding the recurrence equation.
2. Determine the cost of the entire tree by summing the costs of the nodes.

There are other methods for solving recurrence equations; see CLR for details. However, we will use the iteration method almost exclusively.

Example 1: $T(n) = T(n - 1) + 1$


Also derivable: $T(n - 1) = T(n - 2) + 1$
 $T(n - 2) = T(n - 3) + 1$
etc.



Total cost of nodes = number of nodes = n

Example 2: $T(n) = T(n - 1) + n$

Also derivable: $T(n - 1) = T(n - 2) + (n - 1)$
 $T(n - 2) = T(n - 3) + (n - 2)$
 etc.




Total cost of nodes = $1 + 2 + 3 + \dots + (n - 2) + (n - 1) + n$ {Arithmetic series; see below}

$$= \sum_{k=1}^n k =$$

Example 3: $T(n) = T(n/2) + 1$

Also derivable: $T(n/2) = T(n/4) + 1$
 $T(n/4) = T(n/8) + 1$
 etc.




Total cost of nodes = number of nodes = $\lg(n) + 1 = (\lg(n))$

(We start with one $T(n)$ node and generate a number of subnodes equal to the number of times n can be successively divided by 2. The latter quantity is $\lg(n)$; including the 1 generated by the original $T(n)$ node gives $\lg(n) + 1$ nodes.)

Example 4: $T(n) = 2T(n/2) + 1$

Also derivable: $T(n/2) = 2T(n/4) + 1$
 $T(n/4) = 2T(n/8) + 1$
 etc.




Total cost = number of nodes
 = sum of nodes at each level
 = $1 + 2 + 4 + 8 + \dots + 2^{\lg(n)}$ {Geometric series!}

$$= \sum_{k=0}^{\lg(n)} 2^k =$$

Example 5: $T(n) = 2T(n/2) + n$


Also derivable: $T(n/2) = 2T(n/4) + n/2$
 $T(n/4) = 2T(n/8) + n/4$



$$\text{Total cost} = (n)(\text{number of levels}) = n(\lg(n) + 1) = (n \lg(n))$$

Example 6: $T(n) = T(n/2) + n$

Also derivable: $T(n/2) = T(n/4) + n/2$
 $T(n/4) = T(n/8) + n/4$



$$\text{Total cost} = n + n/2 + n/4 + \dots + n/2^{\lg(n)} = \sum_{k=0}^{\lg(n)} \frac{n}{2^k} < \sum_{k=0}^{\lg(n)} \frac{n}{2^k} =$$

Summations

Sigma notation:

$$\sum_{j=1}^n a_j = a_1 + a_2 + \dots + a_n = \sum_{k=1}^n a_k$$

Linearity Laws:


$$\sum_{k=1}^n c a_k = c \sum_{k=1}^n a_k$$

$$\sum_{k=1}^n (a_k + b_k) = \sum_{k=1}^n a_k + \sum_{k=1}^n b_k$$

Arithmetic Series

A series is arithmetic if $a_k = c + a_{(k-1)}$


Trick: Sum corresponding pairs of numbers:



$$\sum_{k=1}^n a_k = \frac{n}{2}(a_1 + a_n)$$

Examples:

- $1 + 2 + 3 + \dots + n =$



- Sum of first n elements of series $7 + 10 + 13 + 16 + \dots$

Geometric Series

A series is geometric if $a_k = c a_{(k-1)}$

Let $S(n)$ stand for
$$\sum_{k=0}^n a_0 c^k = a_0 + a_0 c + a_0 c^2 + a_0 c^3 + \dots + a_0 c^n$$

(Note carefully: $S(n)$ has $n+1$ terms, *not* n terms!)

Notice the following:

$$\begin{aligned} cS(n) &= a_0 c + a_0 c^2 + a_0 c^3 + \dots + a_0 c^n + a_0 c^{n+1} \\ - S(n) &= a_0 + a_0 c + a_0 c^2 + a_0 c^3 + \dots + a_0 c^n \\ \hline (c-1)S(n) &= a_0 c^{n+1} - a_0 = a_0(c^{n+1} - 1) \end{aligned}$$

$$S(n) = \frac{a_0(c^{n+1} - 1)}{(c - 1)}$$

If $0 < c < 1$ and $n \rightarrow \infty$, the above formula can be rewritten as:


$$\lim_{n \rightarrow \infty} S(n) = \frac{a_0}{(1 - c)} \quad \text{if } 0 < c < 1$$

Examples:

$$1 + 2 + 4 + 8 + \dots + 2^n =$$

$$1 + 2 + 4 + 8 + \dots + 2^{\lg(n)} =$$

$$1/2 + 1/4 + 1/8 + \dots =$$



$$1/3 + 1/9 + 1/27 + \dots =$$
